

Heavy Quark Production
and Gluon Shadowing
at RHIC and LHC

Jörg Raufeisen, SLAC

Introduction

Heavy Quark production is one of the most interesting processes to be studied at RHIC and LHC:

- The reaction evolves over large longitudinal distances:

$$G \sim \int |M|^2 d(P.S.)$$

$\hookrightarrow \ln \frac{M^2}{S} \gg 1$ can compensate smallness of α_s

→ pronounced nuclear effects make it possible to study space-time evolution of heavy quark production

- Saturation:

for $M^2 < Q_s^2$, gluon recombination leads to nonlinear terms in the evolution equation:

$$\alpha_s \sim \begin{cases} 1 \text{ GeV at RHIC} \\ 1.5 \text{ GeV at LHC} \end{cases} \quad Q_s^2 \sim A^{1/3}$$


$$\text{new hard scale} \rightarrow \frac{Q_s^2}{M^2} \sim O(1)$$

→ no twist expansion possible

Dipole Approach to Heavy Quark Production

Nikolaev, Piller, Zalicharov, JETP 81 (95) 851

Kopeliovich, Tarasov, hep-ph/0205151
NPA 710 (02) 190



$$\frac{d\sigma_{q\bar{q}}}{dy} = \propto, G(x_1, \mu) \int d\alpha \int d^2 p |\psi_{G \rightarrow q\bar{q}}(\alpha, p)|^2 G_{q\bar{q}G}^N(x_2, \alpha, p)$$

$$\mu \sim m_\alpha$$

General rule:

$$\sigma(aN \rightarrow bcX) = \int d\Gamma |\psi_{a \rightarrow bc}(\Gamma)|^2 \sigma_{bc\bar{c}}^N(\Gamma)$$

Γ : set of all internal variables
of the (bc)-system


$\psi_{a \rightarrow bc}$: LC wavefunctions for
the transition $a \rightarrow bc$

$\sigma_{bc\bar{c}}^N$: cross section for scattering
the (bc \bar{c})-system on a nucleon

The total $c\bar{c}$ -pair cross section

Check validity of dipole approach in $p\bar{p}$
before application to nuclei

(J.R. + J.-C. Peng PRD 67, 054008 (2003))



\hookrightarrow PM calculation from


Nason, Dawson, Ellis, NPB 303 (88) 607

NPB 327 (89) 49

Mangano, Nason, Ridolfi, NPB 373 (92) 295

- Dipole approach describes all data, even at low \sqrt{s} .
- No significant difference between dipole approach and parton model despite $m_c \sim Q_s$

Theoretical uncertainties



Uncertainties arise from the choice of


- i) charm quark mass m_c
- ii) renormalization scale μ_R (enters α_s)
- iii) factorization scale μ_F (enters PDFs)

$$1.2 \text{ GeV} \leq m_c \leq 1.8 \text{ GeV}, \quad m_c \leq \mu_R \leq 2m_c$$

$$\mu_F = 2m_c \text{ fixed}$$

fix parameters to describe data

The total $b\bar{b}$ -pair cross section




$$4.5 \text{ GeV} \leq m_b \leq 5 \text{ GeV} \quad m_b = \mu_C / \mu_F \leq 2 m_b$$

- For $b\bar{b}$ production, one needs the DGLAP improved saturation model of
Bartels et al., PRD 66:014001 (2002)
- The dipole approach tends to predict higher values than the next-to-leading order parton model.

Nuclear Shadowing in Open Charm Production

Two contributions to shadowing:

1) rescattering of $c\bar{c}$ - Fock state



$c\bar{c}$ - separation $\rho \sim \frac{1}{m_c}$

→ higher twist shadowing


$$\text{but } \frac{\Delta G}{G} \sim \frac{A^{1/3} Q_s^2(x)}{m_c^2}$$

not negligible

(Kopeliovich, Tarasov, hep-ph/0205151)
NPB 610 (2001) 160

2) rescattering of higher Fock states

(Gluon Shadowing)



single scattering $c\bar{c}$
sees reduced gluon
density

double scattering of $c\bar{c}G$ Fock state

leads to the leading twist gluon shadowing

Nuclear Shadowing in the Dipole Formulation

Consider very low x_2 : $x_2 \ll 0.1 \text{ fm}^{-1}$

→ $Q\bar{Q}$ pair exists over very large longitudinal distances:

$$l_c \gg R_A$$

→ Partonic configurations with fixed transverse separations are interaction eigenstates:

$$G_{q\bar{q}G}^A = 2 \int d^2 b \left\{ 1 - e^{-\frac{1}{2} \theta_{q\bar{q}G}^P \tilde{T}(b)} \right\}$$


(Eikonalization gives only higher twist shadowing for HQ .)

→ The leading twist gluon shadowing R_G reduces the effective nuclear thickness:

$$\tilde{T}(b) = T(b) R_G(b)$$

Green function technique

Under realistic conditions, the size of the $Q\bar{Q}$ -pair is not completely frozen (finite coherence length)




The lifetime of the $Q\bar{Q}G$ -fluctuation is even shorter.

Describe propagation of $Q\bar{Q}$ (or $Q\bar{Q}G$) through the nucleus by Green function G

Limits: "Eikonal" for $x_2 \rightarrow 0$
PP (single scattering) for $x_2 \rightarrow 0.1$

Nuclear suppression of open charm



Kopeliovich, Tarasov NPA 710 (2002) 180

P_T -Broadening at small and not-so-small x_c

$$\delta \langle P_T^2 \rangle = \langle P_T^2 \rangle_{nA} - \langle P_T^2 \rangle_{nP} \geq 0 \quad \text{for DY, } \gamma/\psi, \gamma$$

- very small x_c : broadening = color filtering
The fluctuation containing the DY or γ/ψ is formed far upstream, but the nucleus filters out large (i.e. small p_T) fluctuations.

⇒ increase of $\langle P_T^2 \rangle$

(Kopeliovich, Raufisen, Tarasov, Johnson,
PRC 67(2003)014903)

- larger $x_c \sim 0.1$:

Broadening is due to multiple rescattering of the projectile parton.


$$\delta \langle P_T^2 \rangle = \underbrace{\frac{\pi^2}{3} \alpha_s (\delta \langle P_T^2 \rangle) \langle F^2 \rangle}_{\substack{\text{Johnson} \\ \text{Kopeliovich, Tarasov,} \\ \text{PRC 63(2001)035203}}} \cdot \langle T \rangle$$

$$C = \left. \frac{d}{d \rho^2} G_{q\bar{q}}(s, \rho^2) \right|_{\rho^2 \rightarrow 0}$$

Mean transverse color field strength:

$$\langle F^2 \rangle = \frac{1}{2\pi p^+} \int dy^- \langle N | F_{\mu}^{+\omega}(y^-) F_{\mu \omega}(0) | N \rangle$$

p_T -broadening at fixed target energies




$$\delta \langle p_T^2 \rangle = \left(\frac{g}{q} \right) G \left\{ \langle T_A \rangle - \langle T_D \rangle \right\}$$

$$G = \left. \frac{d}{d \rho^2} G_{q\bar{q}}(\rho^2, s) \right|_{\rho^2 \rightarrow 0}$$

BDMPS transport coefficient:

$$q_A (\sqrt{s} = 10 \text{ GeV}) \approx 0.08 \frac{\text{GeV}^2}{\text{fm}}$$

$$q_A (\sqrt{s} = 22 \text{ GeV}) \approx 0.1 \frac{\text{GeV}^2}{\text{fm}}$$



Upcoming E866 data
might yield larger
 $\delta \langle p_T^2 \rangle$

Summary

- At high energies, heavy quark production can be formulated in terms of the same color dipole cross section as low- x DIS.
We tested this approach in pp.
- All nuclear effects are then predicted:
no remaining free parameters.
- The dipole approach also includes higher twist contributions to nuclear effects in heavy quark production. (important for charm)
- Nuclear broadening can be calculated in the dipole approach as well:
 - $\delta\langle p_T^2 \rangle$ well reproduced for Υ/ψ and Υ
 - Drell-Yan data overestimated by factor ~2
- The parameter free calculation of $\delta\langle p_T^2 \rangle$ provides an independent estimate of the BDMPS transport coefficient.